

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

(Preservative Free)

MOXIEYE-DTM

MOXIFLOXACIN & DEXAMETHASONE EYE DROPS

COMPOSITION

Moxifloxacin Hydrochloride IP equivalent to Moxifloxacin 0.5% w/v, Dexamethasone Sodium Phosphate IP equivalent to Dexamethasone Phosphate 0.1% w/v, Water for Injections IP q.s.

DESCRIPTION : MOXIEYE-D is a combination of Moxifloxacin 0.5% w/v and Dexamethasone Phosphate 0.1% w/v, Moxifloxacin is a fourth generation fluoroquinolone controls infection by inhibiting the DNA gyrase & topoisomerase IV, Dexamethasone a potent corticosteroid effectively controls the inflammation by inhibiting the release inflammatory mediators.

INDICATIONS: MOXIEYE-D is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

DOSAGE : One or two drops instilled into the conjunctival sac(s), every 4 to 6 hours. During the initial 24 to 48 hours, the dosage may be increased to 1 or 2 drops every two hours. Frequency must be decreased gradually or warranted by improvement in clinical signs. Care should be taken not to discontinue the therapy prematurely.

CONTRAINDICATIONS : MOXIEYE-D is contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis), vaccinia, varicela, and in many other viral diseases of the conjunctiva and cornea. Mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of medication.

WARNING AND PRECAUTIONS : NOT FOR INJECTION.

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision and posterior subcapsular cataract formation.

ADVERSE EFFECTS : The most frequently reported drug-related undesirable effects seen with moxifloxacin are conjunctival irritation, increased lacrimation, keratitis and papillary conjunctivitis.







PACKING : MOXIEYE-D Eye Drops are available in 10ml in *lyondellbasel Purell* low density polyethylene dispensing system consisting of plastic bottle with good flexibility. Tamper evidence ring around the closure and neck area of the bottle.

STORAGE INSTRUCTIONS : Store in cool & dry place. Protect from light.

Keep out of reach of children.

Use the solution within one month after opening the container.

For external use only.

INSTRUCTIONS FOR USE		
 <p>1</p>	 <p>2</p>	 <p>3</p>
<p>Rotate cap anticlockwise to break the seal.</p>	<p>The vial is now ready for use; turn it up side down. Squeeze the walls of the vial gently to deliver sterile drop into the eye.</p>	<p>Replace the cap. Tighten it firmly and keep the vial closed for subsequent use.</p>
 <p>4</p>	 <p>5</p>	 <p>6</p>
<p>Do not touch the nozzle.</p>	<p>Do not rinse the nozzle.</p>	<p>Do not expose to Sunlight.</p>
<p>State of the art technology from Eyekare Ltd.</p>		

Marketed by:



Division of :
Kilitch Healthcare India Ltd.

© Trade Mark applied for

Manufactured in India by:

KH KILITCH HEALTHCARE INDIA LTD.
R-904, 905, T.T.C. Industrial Area, M.I.D.C.,
Navi Mumbai, Dist. Thane, Rabale - 400 701.