

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

TIMOEYETM

TIMOLOL MALEATE EYE DROPS IP 0.5% W/V

COMPOSITION

Timolol Maleate IP equivalent to Timolol 0.5% w/v

Stabilized Oxychloro Complex (as Preservative) ... 0.005% w/v

Purified Water IP q.s.

Other Ingredient: β -Cyclodextrin IP

INTRODUCTION

Timoeye (Timolol Maleate) is a nonspecific beta blocker with no significant intrinsic sympathomimetic activity. Being water soluble and free from local anaesthetic properties, it is suitable for topical administration.

MODE OF ACTION

Timolol maleate is reported to suppress the formation of an aqueous humor and thereby reduce the intraocular pressure. Topical application of Timolol maleate does not affect endothelial permeability.

INDICATIONS

Timoeye (Timolol) is indicated in chronic open angle glaucoma, ocular hypertension, aphakic glaucoma, secondary glaucoma.

ADVERSE EFFECTS

Local irritation, photophobia, blurred vision and itching.

DOSAGE

One drop twice daily or as advised by the Ophthalmologist.

When a patient is transferred from a single anti-glaucoma agent. Continue the drug in use and add a drop of Timoeye in each affected eye twice a day. On the following day, discontinue the formerly used drug completely and continue with Timoeye.

PHARMACEUTICAL PRECAUTIONS

It is desirable that the contents should not be used for more than a month after opening container. Do not touch the dropper tip or let it touch any surface since this may contaminate the solution. If irritation persists or increases, discontinue the use and consult the doctor.

PACKING :

Timoeye Eye Drops are available in 5ml in **lyondellbasel Purell** low density polyethylene dispensing system consisting of plastic bottle with good flexibility. Tamper evidence ring around the closure and neck area of the bottle.







STORAGE :

Store at a temperature not exceeding 30°C. Protect from light and moisture.

Keep out of reach of children.

Use the solution within one month after opening the container.

For external use only.

INSTRUCTIONS FOR USE		
 <p>1</p>	 <p>2</p>	 <p>3</p>
Rotate cap anticlockwise to break the seal.	The vial is now ready for use; turn it up side down. Squeeze the walls of the vial gently to deliver sterile drop into the eye.	Replace the cap. Tighten it firmly and keep the vial closed for subsequent use.
 <p>4</p>	 <p>5</p>	 <p>6</p>
Do not touch the nozzle.	Do not rinse the nozzle.	Do not expose to Sunlight.
State of the art technology from Eyekare Ltd.		

 Trade Mark applied for

Marketed by:



Division of :

Kilitch Healthcare India Ltd.

Manufactured in India by:

KH KILITCH HEALTHCARE INDIA LTD.

R-904, 905, T.T.C. Industrial Area, M.I.D.C.,

Navi Mumbai, Dist. Thane, Rabale - 400 701.

EV032/R01