

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

TROPIEYETM

TROPICAMIDE & PHENYLEPHRINE HYDROCHLORIDE OPHTHALMIC SOLUTION

COMPOSITION :

Tropicamide IP 0.8% w/v, Phenylephrine Hydrochloride IP 5% w/v, Chlorbutol (as Preservative) IP 0.5% w/v, Water for Injections IP q.s.

DESCRIPTION : Phenylephrine/Tropicamide is a combination of a mydriatic/cycloplegic drug. Phenylephrine is a synthetic sympathomimetic (adrenergic mydriatic) following topical application, phenylephrine brings about contraction of the radial muscle thereby bringing about dilatation of the pupil.

Tropicamide is a parasymphatholytic drug which inhibits the contraction of the circular muscle and also paralyses the ciliary muscle. Tropicamide functions as a cycloplegic/mydriatic drug.

INDICATIONS : TROPIEYE can be used both for diagnostic and therapeutic purpose:

- Therapeutic purpose – Inflammatory conditions of the uveal tract
- Diagnostic purpose – Retinal photography – Refractive errors – Fundus examination/photography
- Slit lamp examination
- Pre-operative use – In order to undergo surgical procedure that requires the visualization of structures behind the iris, such as cataract extraction, vitrectomy and retinal detachment surgery, the pupil must be adequately dilated before surgery.

CONTRAINDICATIONS : Hypersensitivity to any of the components. Narrow angles or narrow angle glaucoma.

DOSAGE : Therapeutic purpose Uveitis: 1-2 drops bid-qid or as required.

Diagnostic purpose/Pre-operative purpose: 1-2 drops in the eye(s) 15-30 minutes prior to the procedure.







WARNINGS & PRECAUTIONS : Use with caution when administered with, or up to 21 days after administration of MAO inhibitors as exaggerated adrenergic effects may result. The pressor response of adrenergic agents may also be potentiated by tricyclic antidepressants.

PACKING : Tropieye Eye Drops are available in 5ml in **lyondelbasei Purell** low density polyethylene dispensing system consisting of plastic bottle with good flexibility. Tamper evidence ring around the closure and neck area of the bottle.

STORAGE : Store at a temperature not exceeding 30°C and protect from light. Keep out of reach of children.

Use the solution within one month after opening the container.

For external use only.

INSTRUCTIONS FOR USE		
 <p>1</p>	 <p>2</p>	 <p>3</p>
<p>Rotate cap anticlockwise to break the seal.</p>	<p>The vial is now ready for use; turn it up side down. Squeeze the walls of the vial gently to deliver sterile drop into the eye.</p>	<p>Replace the cap. Tighten it firmly and keep the vial closed for subsequent use.</p>
 <p>4</p>	 <p>5</p>	 <p>6</p>
<p>Do not touch the nozzle.</p>	<p>Do not rinse the nozzle.</p>	<p>Do not expose to Sunlight.</p>
<p>State of the art technology from EyeKare Ltd.</p>		

Marketed by:



Division of:

Kilitch Healthcare India Ltd.

Manufactured in India by:

KH KILITCH HEALTHCARE INDIA LTD.

R-904, 905, T.T.C. Industrial Area, M.I.D.C.,
Rabale, Navi Mumbai - 400 701, Dist. Thane.

TM Trade Mark applied for