

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory



CARBOXYMETHYLCELLULOSE EYE DROPS IP 1% WV

COMPOSITION

Carboxymethylcellulose Sodium IP 1% w/v
Stabilized Oxychloro Complex (as Preservative) 0.0075% w/v
Water for Injections IP q.s.

INTRODUCTION

Weteye Gel Lubricant Eye Drops provides soothing relief for dry, irritated eyes with a formula that resembles your body's own tears. Weteye Gel contains a unique, mild, non-sensitizing preservative that when used, ultimately changes into components of natural tears (sodium chloride + water). Use Weteye Gel for long-lasting relief of your dry, irritated eyes as often as needed.

INDICATIONS

For temporary relief of burning, irritation and discomfort due to dryness of the eye or due to exposure to wind or sun. Also may be used as protectant against further irritation.

DIRECTIONS

Instil 1 or 2 drops in the affected eye(s) as needed. If irritation persists or increases discontinue the use and consult your doctor.

WARNINGS

To avoid contamination, do not touch dropper tip of the container to any surface. Replace cap after using. If you experience eye pain, changes in vision, continued redness or irritation of the eye or if the condition worsens or persists for more than 72 hours, discontinue the use and consult a doctor.

SIDE EFFECTS

When used as directed, no significant side effects are anticipated. Ocular irritation and allergic reactions have been reported occasionally.

PACKING

Weteye Gel are available in 10ml in **lyondellbasel Purell** low density polyethylene dispensing system consisting of plastic bottle with good flexibility. Tamper evidence ring around the closure and neck area of the bottle.

Storage Instructions :







Store in cool & dry place.

Protect from light.

Keep out of reach of children.

Use the solution within one month after opening the container.

For external use only.

INSTRUCTIONS FOR USE		
 1	 2	 3
Rotate cap anticlockwise to break the seal.	The vial is now ready for use; turn it up side down. Squeeze the walls of the vial gently to deliver sterile drop into the eye.	Replace the cap. Tighten it firmly and keep the vial closed for subsequent use.
 4	 5	 6
Do not touch the nozzle.	Do not rinse the nozzle.	Do not expose to Sunlight.
State of the art technology from Eyekare Ltd.		

® Registered Trade Mark

Marketed by:



Division of :
Kilitch Healthcare India Ltd.

Manufactured in India by:

KH KILITCH HEALTHCARE INDIA LTD.
R-904, 905, T.T.C. Industrial Area, M.I.D.C.,
Navi Mumbai, Dist. Thane, Rabale - 400 701.